

SIGNET-RING CELL GASTRIC CARCINOMA IN A DOG

M.M. Noordin, P.K. Low, H.H. Lee, G.K. Dhaliwal and S. Abdul-Jamil

Faculty of Veterinary Medicine
Universiti Putra Malaysia
43400 UPM Serdang, Selangor, Malaysia

SUMMARY

Owing to its rarity, the clinical sign of chronic vomiting is usually diagnosed as foreign body obstruction or as an intestinal accident. Unfortunately, gastric carcinoma is usually never considered as a differential in such cases. This paper describes the pathology of signet-ring cell gastric carcinoma in a dog that had a history of chronic vomiting and haemetemesis along with associated haemogram and clinical biochemical changes pertaining to this tumour.

Keywords: chronic vomiting, haemetemesis, signet-ring cell gastric carcinoma

INTRODUCTION

Although rare in domestic animals, the prevalence of gastric carcinoma is highest in dogs (Withrow, 1996). The stenotic effect of this carcinoma on the stomach is known to lead to vomiting, abdominal pain, anaemia, weight loss, haemetemesis and malaena in dogs (Fonda *et al.*, 1989). In all cases, vomiting is the predominant sign which may appear as early as six months prior to diagnosis. This paper describes a case of gastric carcinoma in a dog with the characteristic clinical-pathological feature of this tumour.

CASE HISTORY

The dog was a 14 year-old spayed Spitz that had a history of chronic vomiting, inappetence and haemetemesis for the past month. On physical examination, the dog was found to be thin, depressed and anaemic. However, no abdominal pain or masses was found. This led to differential diagnoses of gastritis, hepatic or renal insufficiency and neoplasia. Blood was taken for liver and kidney enzymes while treatment for vomiting and inappetence was advocated. Oral metoclopramide (Metoclopramide Omega®) and vitamin B complex were prescribed.

The following day, the dog was brought in again since it continued to vomit despite being given the prescribed drug. Thus, another set of treatment was prescribed with the addition of fluid therapy for 5% dehydration, tagamate (Cimetidine®) and carafate (Sulcrafate®) for gastritis/ulceration and amoxicillin (Amoxil®). As vomiting did not cease, the regime for metoclopramide was changed from oral to the intravenous route.

It was hospitalised for seven days prior to its death on 24 August 2004. During hospitalisation, blood samples were taken. Survey and contrast (Iohexol, Omnipaque®)

radiography and ultrasound were performed on the abdomen.

Clinico-pathologic findings

During hospitalisation, the frequency and severity of vomiting did not subside despite the treatment. The haemogram revealed a regenerative anaemia with a left shift (Table 1) and the vomitus was confirmed to contain blood.

Table 1: Selected blood parameters of the dog following the first day of hospitalisation

Parameter	Reference Values	Result
Erythrocytes (X1012/l)	5.5-8.5	3.43
Haemoglobin (g/l)	74	120-180
PCV (l/l)	0.35-0.55	0.22
Lueocytes (corrected) (X109/l)	6.0-17.0	25.9
Band neutrophils (X109/l)	0.0-0.3	2.33
Segmented neutrophils (X109/l)	3.0-11.5	24.72
Nucleated RBCs (/100WBC)	Rare	2
Reticulocytes (/100RBC)	0.5-1.5	2.4
Plasma protein (g/l)	60-80	72
Icteric index (units)	2-10	2
ALT (U/l)	5-90	25.5
BUN (mmol/l)	3-7.5	6.1
Creatinine (umol/l)	88-176	117

The survey X-ray revealed a lobulated liver while the contrast radiogram showed filling defect at the pyloric region. Thickening of gastric wall along with a lobulated liver was scanned on ultrasound.




Fig. 1. Photograph of the thickened and stenotic pylorus (arrow)




Fig. 4. Photomicrograph of the tumour in the pylorus depicting the presence of signet-ring cell (H&E, X100)



Fig. 2. Photograph of the mottled and shrunken lobulated liver



Fig. 5. Photomicrograph of the tumour in the stomach showing signet-ring cell being laden with mucin (PAS, X400)




Fig. 3. Photomicrograph of the tumour denoting its scirrhus nature in the pylorus (H&E, X 40)




Fig. 6. Photomicrograph of the liver showing clusters of scirrhus (arrows) appearance of the tumour cells

Post mortem findings

Grossly, there was a 4 cm thick greyish scirrhous mass encircling the pyloric mucosa (Fig. 1). This annular fold of mass was found to obliterate more than 85% of the pyloric lumen. Apart from a thickened gastric mucosa, the mottled liver was also nodular and slightly smaller (Fig. 2).

Microscopically, the scirrhous mass consisted of massive fibrous stroma along with sac-like nest of cells (Fig. 3). Although the cells were pleomorphic, the general appearance tended to show up as cuboidal cells, many of which also exhibited signet-ring forms (Fig. 4). The PAS stain performed on all tumour tissues (pylorus and liver) demonstrated that majority of the signet-ring cells were laden with mucin (Fig. 5).

Hepatocytes were in different stages of fatty degeneration, and in some areas, scirrhous arrangements of cells similar to those seen in the mass were evident (Fig. 6). Such cells were also seen to be present in blood vessels and its vicinity.

DISCUSSION

In majority of gastric carcinoma cases, vomiting is the most consistent and predominant clinical signs observed (Fonda *et al.*, 1989). Vomiting can be seen as early as one to six months prior to the diagnosis of this tumour in dogs. In our case, the vomiting was more of haematemesis as confirmed by the presence of occult blood in the vomitus. Owing to the size of the tumour and the frequency of vomiting during hospitalisation, the dog could have been having bouts of vomiting for more than a month as stated.

The radiography and ultrasound results aided the diagnosis of this tumour. Hypoechoic transmural gastric wall thickening with altered layering as found in this dog during ultrasound examination is highly suggestive of gastric carcinoma (Beck *et al.*, 2001). This feature is a result of the tumour arising from the submucosa and massively infiltrating the gastric musculatures (Pennick *et al.*, 1998).

Another factor that might have easily led to the suspicion of gastric carcinoma in this dog is its age. Dogs older than three years are usually affected by gastric carcinoma (Gualtieri *et al.*, 1999) and in our case, the age of 14 years surpassed the mark of age susceptibility to this tumour.

Likewise, the region (pylorus) where the tumour was found is also a predilection site for this tumour. It has been observed that gastric carcinoma commonly occurs at the pyloric, cardia and antral region while the fundus is a rare site. In addition, the gross annular and stenosing (Lingeman *et al.*, 1971; Fonda *et al.*, 1989) appearance of the tumour in the case presented here further confirms the likelihood of a gastric carcinoma.

Undoubtedly, the cellular feature (both on H&E and PAS stains) of the tumour conforms to that of a gastric carcinoma. The pronounced fibrosis, infiltration clusters of pleomorphic cells throughout all layers of the stomach as evident in this case are striking features of this tumour. Furthermore, the presence of signet-ring cells laden with mucin endorsed the definitive diagnosis of gastric carcinoma (Fonda *et al.*, 1989).

The presence of such cells in the liver signified the aggressive nature of this tumour and can be classified as T3N0M1 based on the WHO classification (Hamilton and Aaltonen, 2000). It is also believed that degenerative changes in the liver are possibly secondary to the presence of this tumour due to the competition between tumour cells and hepatocytes for nutrient. Alternatively, compression of the vessels and sinusoids by the tumour cells could also have led to such degenerative changes.

No correlation exists between high prevalence in humans as compared to animals in this area. This could be due to animals being better at avoiding prolonged contact with carcinogens or that animals are inherently more resistant than man to the carcinogens. Thus, the rarity of this case in dogs is simply because Malaysians have a very low rate of gastric carcinoma (Kurioshi *et al.*, 1994).

Several factors are suspected to play a role in gastric carcinogenesis, including the effects of diet, exogenous chemicals, intra-gastric synthesis of carcinogens, genetic factors, infectious agents and pathological conditions in the stomach (Stadtländer and Waterbor, 1999). Likewise, it is almost impossible to determine the exact aetiology in the case presented owing to the prolonged onset of the signs and rather poor history.

Although a number of genetic and molecular alterations have been described (Gao *et al.*, 2001; Tamura *et al.*, 2001; Yokozaki *et al.*, 2001), the pathogenesis of gastric carcinoma remains is still in a state of flux. However, the mechanism by Correa (1992) postulating the initial involvement of gastritis ending in metaplasia, dysplasia, and finally gastric carcinoma is widely accepted. It is very likely that the dog in the case reported could have also been going through such pathogenetic mechanisms. This is because of slow onset or a long period of vomiting.

Finally, the alteration or loss of E-cadherin expression seems to be critical for the development of gastric carcinomas. Furthermore, abnormal E-cadherin-catenin complex expression occurred in early gastric carcinomas and changes in E-cadherin expression might be early events in gastric carcinoma (Blok *et al.*, 1999; Xiangming *et al.*, 1999; Ohene *et al.*, 2000). Abnormal expression of the E-cadherin-catenin complex occurs frequently in gastric carcinoma, and is closely related to its histogenesis (Zhou *et al.*, 2002). In our case, no further immunohistochemical test was done since the microscopic appearance of the tumour stained by H&E and PAS were characteristic of gastric carcinoma.

Thus, the use of scout and contrast radiography, ultrasound, possibly endoscopy (not done in the present study) along with appropriate histochemical examination are essential tools in diagnosing gastric carcinoma in dogs.

REFERENCES

- Beck, C., Slocombe, R.F., O'Neill, T. and Holloway, S.A. (2001). Ultrasound investigation of gastric carcinoma in a dog. *Aust. Vet. J.* **79**: 332-334.
- Blok, P., Craanen, M.E., Dekker, W. and Tytgat, G.N. (1999). Loss of E-cadherin expression in early gastric cancer. *Histopathology* **34**: 410-415.
- Correa, P. (1992). Human gastric carcinogenesis: a multistep and multifactorial process—first American Cancer Society Award Lecture on cancer epidemiology and prevention. *Cancer Res.* **52**: 6735-6740.
- Fonda, D., Gualtieri, M. and Scanziani, E. (1989). Gastric carcinoma in the dog : a clinicopathological study of 11 cases. *J. Small Anim. Pract.* **30**: 353-360.
- Gao H.J., Yu, L.Z., Bai, J.F., Peng, Y.S., Sun, G., Zhao, H.L., Miu, K., Liu, X.Z., Zhang, X.Y. and Zhao, Z.Q. (2001). Multiple genetic alterations and behavior of cellular biology in gastric cancer and other gastric mucosal lesions: H.pylori infection, histological types and staging. *World. J. Gastroenterol.* **6**: 848-854.
- Gualtieri, M., Monzeglio, M.G., and Scanziani, E. (1999). Gastric neoplasia. *Vet. Clin. North Am. Small Anim. Prac.* **29**: 415-420.
- Hamilton, S.R. and Aaltonen, L.A. (2000). WHO Classification of Tumours- Pathology and Genetics of Tumours of Digestive System. Lyon, IARC Press.
- Kurioshi, T., Hayakawa, N., Kurihara, M. and Aoki, K. (1994). Cancer mortality in 33 countries of the world (1953–1987). In: Cancer Mortality and Morbidity Statistics. Tominaga, S., Aoki, K., Fujimoto, I. and Kurihara, M. (Ed.). Gann Monographs on Cancer Research no. 41. CRC Press, Boca Raton, FL, pp. 167–230.
- Lingeman, C.H., Garner, F.M. and Taylor, D.O.N. (1971). Spontaneous gastric adenocarcinomas of dogs: a review. *J. Natl. Cancer Inst.* **47**: 137-153.
- Ohene-Abuakwa, Y., Noda, M., Perenyi, M., Kobayashi, N., Kashima, K., Hattori, T. and Pignatelli, M. (2000). Expression of the E-cadherin/catenin (alpha-, beta-, and gamma-) complex correlates with the macroscopic appearance of early gastric cancer. *J. Pathol.* **192**: 433-439.
- Pennick, D.G., Moore, A.S. and Gliatto, J. (1998). Ultrasonography of canine gastric epithelial neoplasia. *Vet. Radiol. Ultrasound* **39**: 342-348.
- Stadtländer, C.T. and Waterbor, J.W. (2000). Molecular epidemiology, pathogenesis and prevention of gastric cancer. *Carcinogenesis* **20**: 2195-2208.
- Tamura, G., Sato, K., Akiyama, S., Tsuchiya, T., Endoh, Y., Usuba, O., Kimura, W., Nishizuka, S. and Motoyama, T. (2001). Molecular characterization of undifferentiated-type gastric carcinoma. *Lab Invest* **81**: 593-598.
- Withrow, S.J. (1996). Gastric cancer. In: Small Animal Clinical Oncology. Withrow, S.J. and MacEwen, E.G. (Ed.) 2nd ed. W.B. Saunders Co., Philadelphia. pp 244-248.
- Xiangming, C., Hokita, S., Natsugoe, S., Tanabe, G., Baba, M., Takao, S., Kuroshima, K. and Aikou, T. (1999). Co-occurrence of reduced expression of alpha-catenin and overexpression of p53 is a predictor of lymph node metastasis in early gastric cancer. *Oncology* **57**:131-137.
- Yokozaki, H., Yasui, W. and Tahara, E. (2001). Genetic and epigenetic changes in stomach cancer. *Int Rev Cytol.* **204**: 49-95.
- Zhou, Y.N., Xu, C.P., Fang, D.C., Yang, J.M., Han, B., Li, M. and Qiao, L. (2002). Expression of E-cadherin and b-catenin in gastric carcinoma and its correlation with the clinicopathological features and patient survival. *Wrld. J. Gastroenterol.* **8**: 987-993.